



Biomimetic Cortical Nanocircuits: The BioRC Project

Alice C. Parker

NSF Emerging Models of Technology Meeting

July 24, 2008



USC Viterbi
School of Engineering

- Alice Parker, PI and Chongwu Zhou, Co-PI
- Graduate Assistants
 - Chih-Chieh Hsu - CNT circuits and simulation
 - Jonathan Joshi - CMOS circuits and simulation
 - Ko-Chung Tseng - Mathematical models of interconnectivity
 - Chuan Wang - Carbon nanotube fabrication
- Affiliated Students
 - Adi Azar - Neural architecture
 - Khushnood Irani - 3-D circuit visualization
 - Jason Mahvash - analog circuits
 - Numerous directed research students
- Support for this research has been provided by the Viterbi School of Engineering and the WiSE Program at USC and NSF Grant 0726815.



- Complexity:
 - Synaptic mechanisms - excitatory and inhibitory synapses
 - Dendritic computations and dendritic spikes
 - Quantum stochastic behavior of neurotransmitter release
- Scale:
 - 100×10^9 neurons
 - 10^4 to 10^5 synapses/neuron
 - ~100 transistors/synapse including dendritic computations
 - CMOS neurons for a cortex, absent interconnection area, could occupy an entire room, even in 2021
- Connectivity:
 - Fan-in/neuron 10^4 to 10^5 distinct connections
 - Fan-out 10^4
 - Address space 37 bits (assuming synaptic inputs are distinct)
- Plasticity:
 - New neural connections form within hours
 - Presynaptic depression/facilitation occur
 - Postsynaptic depression and potentiation occur







- Complexity:
- **Exploit the analog computational power of transistor circuits**
- Scale:
- **Consider nanotechnological solutions - nanotubes, nanowires, graphene, quantum dots**
- Connectivity:
 - **3-D structure probably required**
- Plasticity:
- **Add transistors as “knobs” to control neural behavior**
- **Self-assembly, using a protein gel to provide scaffolding, and synthetic DNA to assemble/reconfigure neural circuits**

- **We are very far from a synthetic human cortex, but it may be possible in the coming decades**




- **Carbon nanotube fabrication (Chongwu Zhou)**
 - Aligned nanotubes, logic gates







Biomimetic Neural Circuits





Excitatory synapse



Axon Hillock



Dendritic Tree



Inhibitory Synapse






Figure from Principles of Neural Science [2] p.22

The whole neuron can be divided into these sub-circuits:


- Synapse
 - Excitatory/Inhibitory synapse circuit (Action Potential as inputs and EPSP/IPSP as outputs)
- Dendritic Tree
 - A pool of voltage adders (which can add two input stimuli in both linear or non-linear ways)
- Axon Hillock
 - Amplifier (in order to reach the threshold of carbon nanotube FET)
 - Spike-initiator (Action Potentials are all-or-none)




Results to Date: A Carbon Nanotube Synapse




Results to Date: A Carbon Nanotube Synapse




Results to Date: A CMOS Inhibitory Synapse



Wave	Symbol
D0:tr3:v(ap1)	X - -
D0:tr3:v(out)	* - -



Wave	Symbol
D0:tr1:v(out)	X
D0:tr2:v(out)	○
D0:tr3:v(out)	△
D0:tr4:v(out)	□
D0:tr5:v(out)	◇
D0:tr6:v(out)	*
D0:tr7:v(out)	+



Dendritic Computations

Linear or Non-linear summation


- Schiller et al. compared the measured and arithmetic results of EPSP summation at soma of layer-5 pyramidal neuron with respect to within-branch and between-branch stimulations
- It appears that between-branch EPSP summation is linear for weak and medium stimuli and slightly superlinear for strong stimuli.
- On the other hand, within-branch EPSP summation shows both linearity and non-linearity depending on the strength of EPSP. It was linear – weak EPSP ($\sim <1\text{mV}$), superlinear – medium EPSP ($1\sim 3\text{mV}$), sublinear – strong EPSP ($3\sim 10\text{mV}$)

Adder structure


- Adding two inputs linearly, sublinearly, and superlinearly

Within-branch

Between-branch




A and B are 20 μm separated





— Arithmetic
— Measured at soma




Weak inputs




Medium inputs




Strong inputs





- Shunting Inhibition
 - Pulls the EPSP or the AP down to zero volts
- Hyperpolarizing Inhibition
 - More of a subtractive behavior



Wave	Symbol
D0.tr0:v(epsp)	X
D0.tr0:v(ipsp)	○
D0.tr0:v(dendrite_out)	△



Simplified Central Neuron Circuit



Red: Action Potential
(artificial input to the pre-synaptic terminal)

Green: EPSP from the dendrites (post-synaptic sites) of the neuron

Blue: Action Potential spike
(initiated at the axon hillock of the neuron)

BioRC

The End

USC Viterbi
School of Engineering

Thank You

